

REFERENCES

- Aguilera-Aleorra M. G. (2015). Identification and typing methods for the study of bacterial infections: a brief review and Mycobacterial as case study. *Archives of Clinical Microbiology*, 7:1.
- Ahmadian, A., Ehn, M., & Hober, S. (2006). Pyrosequencing: history, biochemistry and future. *Clinica chimica acta*, 363(1-2), 83-94.
- Al-Haj, Mashan, N. I., Shamsudin, M. N., Mohamad, H., Vairappan, C. S., & Sekawi, Z. (2010). Antibacterial activity of marine source extracts against multidrug resistance organisms. *American Journal of Pharmacology and Toxicology*, 5(2), 95-102.
- Alipiah, N. M., Ramli, N. H. S., Low, C. F., Shamsudin, M. N., & Yusoff, F. M. (2016). Protective effects of sea cucumber surface-associated bacteria against *Vibrio harveyi* in brown-marbled grouper fingerlings. *Aquaculture Environment Interactions*, 8, 147-155.
- Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W., et al. (1997). Gapped BLAST and PSI-BLAST : a new generation of protein database search programs. *Nucleic Acids Res*, 25, 3389–3402.
- Alshamaa, S., & M Issam, M. (2017). The Role of Extracted Carotenoid from Staphylococci as Antioxidant and Antibacterial. *Rafidain Journal of Science*, 26(1), 17-24.
- Anderson, S. C., Flemming, J. M., Watson, R., & Lotze, H. K. (2011). Serial exploitation of global sea cucumber fisheries. *Fish and Fisheries*, 12(3), 317-339.
- Anwar, M. A., & Choi, S. (2014). Gram-negative marine bacteria: Structural features of lipopolysaccharides and their relevance for economically important diseases. *Marine drugs*, 12(5), 2485-2514.

- Amaro, T., Witte, H., Herndl, G., Cunha, M., & Billett. (2009). Deep-sea bacterial communities in sediments and guts of deposit-feeding holothurians in Portuguese canyons (NE Atlantic). *Deep Sea Research Part I Oceanographic Research Papers*, 56, 1834-1843.
- Amaro, T., Luna, G. M., Danovaro, R., Billett, D. S. M., and Cunha, M. R. (2012). High prokaryotic biodiversity associated with gut contents of the holothurian *Molpadia musculus* from the Nazaré Canyon (NE Atlantic). *Deep Sea Research Part I*, 63, 82–90.
- Armstrong, E., Yan, L., Boyd, K. G., Wright, P. C., & Burgess, J. G. (2001). The symbiotic role of marine microbes on living surfaces. *Hydrobiologica*, 461, 37–40.
- Aruldass, C. A., Dufossé, L., & Ahmad, W. A. (2018). Current perspective of yellowish-orange pigments from microorganisms-a review. *Journal of Cleaner Production*, 180, 168-182.
- Avendaño-Herrera, R., Lody, M., & Riquelme, C. E. (2005). Producción de sustancias inhibitorias entre bacterias de biopelículas en substratos marinos. *Revista de Biología Marina y Oceanografía*, 40, 117-125.
- Azam, F., & Worden, A. Z. (2004). Microbes, Molecules, and Marine Ecosystems. *Science*, 303(5664), 1622+.
- Azman, A. S., Mawang, C. I., & Abubakar, S. (2018). Bacterial pigments: the bioactivities and as an alternative for therapeutic applications. *Natural Product Communications*, 13(12).
- Babalola, O. O. (2003). Molecular techniques: An overview of methods for the detection of bacteria. *African Journal of Biotechnology*, 2 (12), 710-713.
- Baccetti, B., & Rosati, F. (1968) The fine structure of polian vesicles of holothurians. *Z Zellforsch Mikrosk Anat*, 90(1):148-60.
- Backhuys, W. (1977). Handbook of the Echinoderms of the British Isles. Uitgever, Rotterdam, Netherlands. Reprint of 1927 edition published by Oxford University Press, Oxford.

Baine, M., & Sze, C. P. (1998). Sea cucumber fisheries in Malaysia, towards a conservation strategy. *SPC Beche-de-mer Information Bulletin*, 12, 6-10.

Bakus, G. J. (1973). The biology and ecology of tropical holothurians. In: Q.A. Jones and R. Endean (eds). *Biology and geology of coral reefs*. Vol. I. Academic Press, London.

Balasingham, A. (2008). *Development of Multi Locus Variable Number Tandem Repeat Analysis:(MLVA) for the genotyping of Legionella Pneumophila isolated from various habitats* (Master's thesis).

Balraj, J., Pannerselvam, K., & Jayaraman, K. (2014). Isolation of pigmented marine bacteria *Exiguobacterium* sp. from the Peninsular Region of India and a study on biological activity of purified pigment. *International Journal of Scientific & Technology Research*, 3(3), 375-384.

Barghouthi, S. A. (2011). A universal method for the identification of bacteria based on general PCR primers. *Indian journal of microbiology*, 51(4), 430-444.

Barretto, D. A., & Vootla, S. K. (2018). In Vitro Anticancer Activity of Staphyloxanthin Pigment Extracted from *Staphylococcus gallinarum* KX912244, a Gut Microbe of *Bombyx mori*. *Indian journal of microbiology*, 58(2), 146-158.

Becker, P. T., & Flammang, P. (2010). Unravelling the sticky threads of sea cucumbers-A comparative study on Cuvierian tubule morphology and histochemistry. In *Biological Adhesive Systems* (pp. 87-98). Springer, Vienna.

Becker, K., Heilmann, C., & Peters, G. (2014). Coagulase-negative staphylococci. *Clinical microbiology reviews*, 27(4), 870-926.

Beilstein, F., & Dreiseikelmann, B. (2006). Bacteriophages of freshwater *Brevundimonas vesicularis* isolates. *Research in Microbiology*, 157(3), 213-219.

Ben-Dov, E., Shapiro, O. H., Siboni, N., & Kushmaro, A. (2006). Advantage of using inosine at the 3' termini of 16S rRNA gene universal primers for the study of microbial diversity. *Applied and Environmental Microbiology*, 72(11), 6902-6906.

Benson, D. A., Karsch-Mizrachi, I., Lipman, D. J., Ostell, J., & Wheeler, D. L. (2004). GenBank: update. *Nucleic acids research*, 32(suppl_1), D23-D26.

Bergeron, M. G., & Ouellette, M. (1998). Preventing antibiotic resistance through rapid genotypic identification of bacteria and of their antibiotic resistance genes in the clinical microbiology laboratory. *Journal of clinical microbiology*, 36(8), 2169-2172.

Berry, A., Janssens, D., Hümbelin, M., Jore, J. P., Hoste, B., Cleenwerck, I., Vancanneyt, M., Bretzel, W., Mayer, A. F., Lopez-Ulibarri, R. & Pasamontes, L. (2003). *Paracoccus zeaxanthinifaciens* sp. nov., a zeaxanthin-producing bacterium. *International Journal of Systematic and Evolutionary Microbiology*, 53(1), 231-238.

Bhat, S., & Amin, T. (2013). Isolation and Characterization of Pigment Producing Bacteria from Various foods for their possible use as biocolours. *International Journal of Recent Scientific Research*, 4(10): 1605-1609.

Bhakuni, D. S., & Rawat, D. S. (2006). *Bioactive marine natural products*. Springer Science & Business Media.

Bibi, F., Yasir, M., Al-Sofyani, A., Naseer, M. I., & Azhar, E. I. (2020). Antimicrobial activity of bacteria from marine sponge *Suberea mollis* and bioactive metabolites of *Vibrio* sp. EA348. *Saudi journal of biological sciences*, 27(4), 1139-1147.

Blockley, A., Elliott, D. R., Roberts, A. P., & Sweet, M. (2017). Symbiotic microbes from marine invertebrates: driving a new era of natural product drug discovery. *Diversity*, 9(4), 49.

Bogatyrenko, E. A., & Buzoleva, L. S. (2016). Characterization of the gut bacterial community of the Japanese sea cucumber *Apostichopus japonicus*. *Microbiology*, 85(1), 116-123.

Bordbar, S., Anwar, F., & Saari, N. (2011). High-value components and bioactives from sea cucumbers for functional foods – a review. *Marine drugs*, 9(10), 1761-1805.

- Brady, S. F., Simmons, L., Kima, J. H., & Schmidt, E. W. (2009). Metagenomic approaches to natural products from free-living and symbiotic organisms. *Natural Product Reports*, 26, 1488–1503.
- Brandt, J.F. (1835). Echinodermata ordo Holothurina. Pp. 42-62. In *Prodromus Descriptionis Animalium Ab. H. Mertensio In Orbis Terrarum Circumnavigatione Observatorum*. Fasc. I: 75 pps. Petropoli.
- Bruckner, A. W., Johnson, K. A., & Field, J. D. (2003). Conservation strategies for sea cucumbers: Can a CITES Appendix II listing promote sustainable international trade?. *SPC Beche-de-mer Information Bulletin*, 18, 24-33.
- Bruhn, J. B., Gram, L., & Belas, R. (2007). Production of antibacterial compounds and biofilm formation by *Roseobacter* species are influenced by culture conditions. *Applied and Environmental Microbiology*, 73, 442-450.
- Bultel-Ponce, V., Berge, J-P., Debitus, C., Nicolas, J-L., & Guyot, M. (1999). Metabolites from the sponge-associated bacterium *Pseudomonas* species. *Marine Biotechnology*, 1, 384-390.
- Bunnoy, A., Na-Nakorn, U., Kayansamruaj, P., & Srisapoome, P. (2019). *Acinetobacter* strain KUO11TH, a unique organism related to *Acinetobacter pittii* and isolated from the skin mucus of healthy bighead catfish and its efficacy against several fish pathogens. *Microorganisms*, 7(11), 549.
- Burgess, J. G., Boyd, K. G., Armstrong, E., Jiang, Z., Yan, L., & Berggren, M. (2003). The development of a marine natural product-based antifouling paint. *Biofouling*, 19, 197–205.
- Campanale, C., Massarelli, C., Savino, I., Locaputo, V., & Uricchio, V. F. (2020). A detailed review study on potential effects of microplastics and additives of concern on human health. *International journal of environmental research and public health*, 17(4), 1212.
- Celedon, R. S., & Diaz, L. B. (2021). Natural pigments of bacterial origin and their possible biomedical applications. *Microorganisms*, 9(4), 739.
- Chanderan, M., Mazlan, N., Weinheimer, A. Z., Young, T. T., & Anua, S. M. (2019). Screening of vibrio spp. isolated from sea cucumber *Stichopus horrens*. *International Journal of Medical Toxicology & Legal Medicine*, 22(1and2), 130-137.

- Chen, L., Wang, X-Y., Liu, R-Z., & Wang, G-Y. (2021). Culturable microorganisms associated with sea cucumbers and microbial natural products. *Marine Drugs*, 19(8), 461.
- Chimetto, L. A., Cleenwerck, I., Moreira, A. P., Brocchi, M., Willems, A., De Vos, P., & Thompson, F. L. (2011). *Vibrio variabilis* sp. nov. and *V. maritimus* sp. nov., isolated from *Palythoa caribaeorum*. *International Journal of Systematic and Evolutionary Microbiology*, 61, 3009-3305.
- Choo, P. S. (2004). Fisheries, trade and utilization of sea cucumbers in Malaysia.
- Choo, P. S. (2008). Population status, fisheries and trade of sea cucumbers in Asia.
- Clauditz, A., Resch, A., Wieland, K. P., Peschel, A., & Gotz, F. (2006). Staphyloxanthin plays a role in the fitness of *Staphylococcus aureus* and its ability to cope with oxidative stress. *Infection and immunity*, 74(8), 4950-4953.
- Conand, C. (2004). Present status of world sea cucumber resources and utilization: An international overview. p. 13–25. In: Lovatelli A., Conand C. Purcell S., Uthicke S., Hamel J.-F. and Mercier A. (eds). *Advances in sea cucumber aquaculture and management*. FAO Fisheries Technical Reports No. 463, FAO, Rome. 425 p.
- Conand, C. (2004). Sea cucumber biology, taxonomy, distribution: conservation Status. In *Proceeding of the Convention on International Trade in Endangered Species of Wild Fauna and Flora. Tech. Workshop on the Conservation of Sea Cucumbers in the Families*.
- Conand, C., Polidoro, B., Mercier, A., Gamboa, R., Hamel, J. F., & Purcell, S. (2014). The IUCN Red List assessment of aspidochirotid sea cucumbers and its implications. *SPC Beche-de-mer Information Bulletin*, 34(5), 3-7.
- Crapart, S., Fardeau, M. L., Cayol, J. L., Thomas, P., Sery, C., Ollivier, B., & Combet-Blanc, Y. (2007). *Exiguobacterium profundum* sp. nov., a moderately thermophilic, lactic acid-producing bacterium isolated from a deep-sea hydrothermal vent. *International journal of systematic and evolutionary microbiology*, 57(2), 287-292.

D'Auria, G., Pushker, R., & Rodriguez-Valera, F. (2006). IWoCS: analyzing ribosomal intergenic transcribed spacers configuration and taxonomic relationships. *Bioinformatics*, 22(5), 527-531.

Darabpour, E., Ardakani, M. R., Motamedi, H., Ghezelbash, G., & Ronagh, M. T. (2010). Isolation of an antibiotic producer *Pseudomonas* sp. from the Persian Gulf. *Asian Pacific Journal of Tropical Medicine*, 3(4), 318-321.

Darwesh, O. M., Barakat, K. M., Mattar, M. Z., Sabae, S. Z., & Hassan, S. H. (2019). Production of antimicrobial blue green pigment Pyocyanin by marine *Pseudomonas aeruginosa*. *Biointerface Res. Appl. Chem*, 9, 4334-4339.

Das, S., Lyla, P., & Khan, S. (2006). Marine microbial diversity and ecology: Importance and future perspectives. *Current Science*, 90(10), 1325-1335.

Divya, J., Mohandas, A., & Singh, B. (2018). A non-pathogenic environmental isolate of *Pseudomonas aeruginosa* MCCB 123 with biotechnological potential. *International Journal of Current Microbiology Applied Sciences*, 7, 3060-3071.

Du, H., Bao, Z., Hou, R., Wang, S., Su, H., Yan, J., Tian, M., Li, Y., Wei, W., Lu, W. & Hu, J. (2012). Transcriptome sequencing and characterization for the sea cucumber *Apostichopus japonicus* (Selenka, 1867). *PloS one*, 7(3), e33311.

Dufossé, L. (2009). Pigments. *Microbial*, 10.

Durán, N., Teixeira M. F. S., de Conti R., & Esposito, E. (2002). Ecological-Friendly Pigments from Fungi. *Critical Reviews in Food Science and Nutrition*, 42(1), 53-66.

Eissa, S. H., Omran, N. E., Salem, H. K., Kabbash, A. K. & Kandeil, M. A. (2017). Surveillance study on the most common sea-cucumbers in some Egyptian coasts. *The Egyptian Journal of Experimental Biology*, 13(2), 300-308.

Elhadi, N., Radu, S., Chen, C. H., & Nishibuchi, M. (2004). Prevalence of potentially pathogenic *Vibrio* species in the seafood marketed in Malaysia. *Journal of Food Protection*, 67(7), 1469-1475.

El-Fouly, M. Z., Sharaf, A. M., Shahin, A. A. M., El-Bialy, H. A., & Omara, A. M. A. (2015). Biosynthesis of pyocyanin pigment by *Pseudomonas aeruginosa*. *Journal of Radiation Research and Applied Sciences*, 8(1), 36-48.

El Samak, M., Solyman, S. M., & Hanora, A. (2018). Antimicrobial activity of bacteria isolated from Red Sea marine invertebrates. *Biotechnology Reports*, 19, e00275.

Enomoto, M., Nakagawa, S. & Sawabe, T. (2012). Microbial communities associated with holothurians: Presence of unique bacteria in the coelomic fluid. *Microbes and Environments*, 27(3), 300-305.

Farjami, B., Nematollahi, M. A., Moradi, Y., Irajian, G. R., Nazemi, M., Ardebili, A., & Pournajaf, A. (2013). Antibacterial activity of the sea cucumber *Holothuria leucospilota*.

Farouk, A. E., Ghouse, F. A. H., & Ridzwan, B. H. (2007). New bacterial species isolated from Malaysian sea cucumbers with optimized secreted antibacterial activity. *American Journal of Biochemistry and Biotechnology*, 3(2), 60–65.

Faulkner, D. J. (2002). Marine natural products. *Natural Product Reports*, 19: 1-48.

Feghali, P. A. R., & Nawas, T. (2018). Pyocyanin: A Powerful Inhibitor Of bacterial Growth and Biofilm Formation. *Inter. J. Case Rep. Stud*, 2(2), 101-107.

Felsenstein, J. (1985). Confidence Limits on Phylogenies: An Approach Using the Bootstrap. *Evolution*, 39(4), 783-791.

Felske, A., Wolterink, A., Van Lis, R., & Akkermans, A. D. (1998). Phylogeny of the main bacterial 16S rRNA sequences in Drentse A grassland soils (The Netherlands). *Applied and Environmental Microbiology*, 64(3), 871-879.

Fitri, D. S., Pangastuti, A., Susilowati, A., & Sutarno, S. (2017). Endophytic bacteria producing antibacterial against methicillin resistant *Staphylococcus aureus* (MRSA)

- in seagrass from Rote Ndao, East Nusa Tenggara, Indonesia. *Biodiversitas Journal of Biological Diversity*, 18(2), 733-740.
- Flammang, P. (2003). The glue of sea cucumber Cuvierian tubules: a novel marine bioadhesive. Ifremer, Plouzane (France).
- Francis, F. J. (2000). Carotenoids as Food Colorants. *Cereal Food World*, 45: 198-203.
- Fredalina, B. D., Ridzwan, B. H., Zainal, A. A., Kaswandi, M. A., Zaiton, H., Zali, I., Kittakoop, P. & Mat Jais, A. A. (1999). Fatty acid compositions in local sea cucumber, *Stichopus chloronotus*, for wound healing. *General Pharmacology*, 33(4): 337-340.
- Gao, F., Li, F., Tan, J., Yan, J., & Sun, H. (2014). Bacterial community composition in the gut content and ambient sediment of sea cucumber *Apostichopus japonicus* revealed by 16S rRNA gene pyrosequencing. *PloS one*, 9(6), e100092.
- Gao, M. L., Hou, H. M., Zhang, G. L., Liu, Y., & Sun, L. M. (2017). Bacterial diversity in the intestine of sea cucumber *Stichopus japonicus*. *Iranian Journal of Fisheries Sciences*, 16(1), 318-325.
- Gaston, K. J., & Spicer, J. I. (2004). *Biodiversity: an introduction*. 2nd Edition. Blackwell
- Glöckner, F.O., Stal, L.J., Sandaa, R. A., Gasol, J.M., O’Gara, F., Hernandez, F., Labrenz, M., Stoica, E., Varela, M. M., Bordalo, A., & Pitta, P. (2012). Marine microbial diversity and its role in ecosystem functioning and environmental change. In: Calewaert JB, McDonough N (eds) Marine Board Position Paper 17. Marine Board-ESF, Ostend, Belgium, pp 13–25
- Godinho, A., & Bhosle, S. (2008). Carotenes produced by alkaliphilic orange-pigmented strain of *Microbacterium arborescens*-AGSB isolated from coastal sand dunes.
- Goecke, F., Labes, A., Wiese, J., & Imhoff, J. F. (2010). Chemical interactions between marine macroalgae and bacteria. *Marine Ecology Progress Series*, 409, 267–300.

- Gould, I. M. (2000). Towards a common susceptibility testing method?. *Journal of Antimicrobial Chemotherapy*, 45(6), 757-762.
- Guo, Y., Ding, Y., Xu, F., Liu, B., Kou, Z., Xiao, W., & Zhu, J. (2015). Systems pharmacology-based drug discovery for marine resources: An example using sea cucumber (Holothurians). *Journal of Ethnopharmacology*, 165, 61-72.
- Gupta, C., Amar, P. G., Prakash, D., Goyal, S., & Gupta, S. (2011). Microbes as potential source of biocolors. *Pharmacologyonline*, 2, 1309–1318.
- Hadfield, M. G. (2011). Biofilms and marine invertebrate larvae: What bacteria produce that larvae use to choose settlement sites. *Annual Review of Marine Science*, 3, 453–470.
- Hamel, J. -F., & Mercier, A. (2000). Cuvierian tubules in tropical holothurians: Usefulness and efficiency as a defence mechanism. *Marine and Freshwater Behaviour and Physiology*, 33, 115-139.
- Hashim, R. B. (1993). *Sumber Makanan Persisiran Laut Sabah*. Kuala Lumpur: Dewan Bahasa dan Pustaka. p. 309.
- Hassani, H. H., Hasan, H. M., Al-Saadi, A., Ali, A. M., & Muhammad, M. H. (2012). A comparative study on cytotoxicity and apoptotic activity of pyocyanin produced by wild type and mutant strains of *Pseudomonas aeruginosa*. *Eur J Exp Biol*, 2(5), 1389-1394.
- Hawa, I., Zulaikah, M., Mohamed, J., AA, Z. A., Kaswandi, M. A., & Ridzwan, B. H. (1999). The potential of the coelomic fluid in sea cucumber as an antioxidant. *Malaysian Journal of Nutrition*, 5(1 & 2), 55-59.
- Health Protection Agency. (2007). Identification of *Staphylococcus* species, *Micrococcus* species and *Rothia* species. National Standard Method BSOP ID 7.
- Hedges, A. J. (1999). The influence of factors affecting the 'critical population' density of inocula on the determination of bacterial susceptibility to antibiotics by disc diffusion methods. *Journal of Antimicrobial Chemotherapy*, 43(2), 313-313.

Henriques, M., Silva, A., & Rocha, J. (2007). Extraction and quantification of pigments from a marine microalga: a simple and reproducible method. *Communicating Current Research and Educational Topics and Trends in Applied Microbiology Formatex*, 2, 586-593.

Hess, M., Sczyrba, A., Egan, R., Kim, T. W., Chokhawala, H., Schroth, G., Luo, S., Clark, D. S., Chen, F., Zhang, T., Mackie, R. I., Pennacchio, L. A., Tringe, S. G., Visel, A., Woyke, T., Wang, Z., & Rubin, E. M. (2011). Metagenomic discovery of biomass-degrading genes and genomes from cow rumen. *Science*, 331, 463-467.

Higgins, M. Sea cucumbers in a deep pickle. Environmental News Network, 30 August 2000.

Hill, G. E., Hill, G. E., McGraw, K. J., & Kevin, J. (Eds.). (2006). *Bird coloration: function and evolution* (Vol. 2). Harvard University Press.

Ho, S. L. (1992). Coral reefs of Malaysia. Tropical Press Sdn. Bhd., Kuala Lumpur, pp. 3-11, 117-119.

Holmström, C., Egan, S., Franks, A., McCloy, S., & Kjelleberg, S. (2002). Antifouling activities expressed by marine surface associated *Pseudoalteromonas* species. *FEMS Microbiology Ecology*, 41(1), 47-58.

Hui, Y. H., & Sherkat, F. (2005). Handbook of Food Science, Technology and Engineering, 4th Ed. London, New York: CRC Press.

Husin, M. J. M., Mazlan, N., Shalom, J., Saud, S. N., & Sani, M. S. A. (2021). Evaluation of Microplastics Ingested by Sea Cucumber *Stichopus Horrens* in Pulau Pangkor, Perak, Malaysia.

Iida, T., Park, K. S., & Honda, T. (2006). *Vibrio parahaemolyticus*. *The Biology of Virus*, 340-348.

Inouye, B. (n.d.). Internal anatomy of a sea cucumber [Illustration]. https://manoa.hawaii.edu/sealarning/media_colorbox/3381/media_original/en

Ismail, H., Punithamalar, A., Sakian, N. I. M., & Hashim, R. (2004). Heavy metal concentration in body walls of Malaysian sea cucumbers. *Jurnal Sains Kesihatan Malaysia (Malaysian Journal of Health Sciences)*, 2(1).

Isnansetyo, A., & Kamei, Y. (2009). Bioactive substances produced by marine isolates of *Pseudomonas*. *Journal of Industrial Microbiology and Biotechnology*, 36(10), 1239-1248.

Jaeger, G. F. (1833). *De Holothuriis. Turici*, 40.

Jafarzade, M., Yahya, N. A., Mohamad, S., Usup, G., & Ahmad, A. (2013). Isolation and characterization of pigmented bacteria showing antimicrobial activity from Malaysian marine environment. *Malaysian Journal of Microbiology*, 9(2), 152-160.

Janda, J., & Abbott, S. (2007). 16S rRNA Gene Sequencing for Bacterial Identification in the Diagnostic Laboratory: Pluses, Perils, and Pitfalls. *Journal of clinical microbiology*, 45, 2761.

Jensen, P. R., & Fenical, W. (1994). Strategies for the discovery of secondary metabolites from marine bacteria: ecological perspectives. *Annual Review of Microbiology*, 48, 559-584.

Jiang, Y., Yao, L., Li, F., Zhijun, T., Yuxiu, Z., & Wang, L. (2014). Characterization of antimicrobial resistance of *Vibrio parahaemolyticus* from cultured sea cucumbers (*Apostichopus japonicas*). *Letters in Applied Microbiology*, 59(2).

Joint, I., Mühling, M., & Querellou, J. (2010). Culturing marine bacteria—an essential prerequisite for biodiscovery. *Microbial biotechnology*, 3(5), 564-575.

Kamarudin, K. R., Rehan, A. M., Lukman, A. L., Ahmad, H. F., Anua, M. H., Nordin, N. F., Hashim, R., Hussin, R., & Usup, G. (2009). Coral reef sea cucumbers in Malaysia. *Malaysian Journal of Science*, 28, 171-186.

Kamarudin, K. R., Hashim, R., & Usup, G. (2010). Phylogeny of sea cucumber (Echinodermata: Holothuroidea) as inferred from 16S mitochondrial rRNA gene sequences. *Sains Malaysiana*, 39(2), 209-218.

Kamarudin, K. R., Rehan, A. M., Hashim, R., Usup, G., Ahmad, H. F., Anua, M. H., & Idris, M. Y. (2011). Molecular phylogeny of *Holothuria* (Mertensiothuria) *leucospilota* (Brandt 1835) as inferred from Cytochrome C Oxidase I Mitochondrial DNA gene sequences. *Sains Malaysiana*, 40(2), 125-133.

Kamarudin, K. R., Ngah, N., Hamid, T. H. T. A., & Susanti, D. (2013). Isolation of a Pigment-producing Strain of *Staphylococcus kloosii* from the Respiratory Tree of *Holothuria* (Mertensiothuria) *leucospilota* from Malaysian Waters. *Tropical life sciences research*, 24(1), 85.

Kamarudin, K. R., & Rehan, M. M. (2015). Morphological and molecular identification of *Holothuria* (Merthensiothuria) *leucospilota* and *Stichopus horrens* from Pangkor Island, Malaysia. *Tropical Life Sciences Research*, 26(1), 87.

Kamarudin, K. R., Usup, G., Hashim, R., & Rehan, M. M. (2015). Sea Cucumber (Echinodermata: Holothuroidea) Species Richness at Selected Localities in Malaysia. *Pertanika Journal of Tropical Agricultural Science*, 38(1).

Kamarudin, K. R. (2018). *Species Identification, Molecular Phylogenetic and Genetic Diversity of Sea Cucumber (Phylum Echinodermata: Class Holothuroidea) in Malaysia* (Doctoral thesis, Universiti Sains Islam Malaysia).

Kerr, A. M., Netchy, K., & Gawel, A. M. (2006). Survey of the shallow-water sea cucumbers of the central Philippines. *A Report to the Municipalities of Negros Oriental, Cebu and Bohol, local Bantay Dagat groups, Coastal Conservation and Education Foundation, Inc., and Silliman University-Angelo King Center for Research and Environmental Management. University Of Guam Marine Laboratory.*

Khartini, A. W., Dayang Fredalina, B., Kaswandi, A., Hing, H. L., Oon, K. W. & Sit, N. W. (2003). Antimicrobial and cytotoxic activities of *Holothuria* sp. and *B. marmorata vitiensis*. *Malaysian Journal of Biochemistry and Molecular Biology*, 8: 49-53.

Khusro, A., Aarti, C., Dusthacker, A., & Agastian, P. (2018). Anti-tubercular and probiotic properties of coagulase-negative staphylococci isolated from Koozh, a traditional fermented food of South India. *Microbial pathogenesis*, 114, 239-250.

Kjørboe, T., Tang, K., Grossart, H. P., & Ploug, H. (2003). Dynamics of microbial communities on marine snow aggregates: colonization, growth, detachment, and grazing mortality of attached bacteria. *Applied and Environmental Microbiology*, 69(6), 3036-3047.

Kim, S-K., & Himaya, S. W. A. (2012). Triterpene glycosides from sea cucumbers and their biological activities. *Advances in Food and Nutrition Research*, 65, 297-319.

Kim, T. Y., Lee, J. J., Kim, B. S., & Choi, S. H. (2017). Whole-body microbiota of sea cucumber (*Apostichopus japonicus*) from South Korea for improved seafood management. *Journal of microbiology and biotechnology*, 27(10), 1753-1762.

Kimura, M. (1980). A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, 16(2), 111-120.

Kleinegris, D. M., van Es, M. A., Janssen, M., Brandenburg, W. A., & Wijffels, R. H. (2010). Carotenoid fluorescence in *Dunaliella salina*. *Journal of applied phycology*, 22(5), 645-649.

Konig, G. M., Kehraus, S., Seibert, S. F., Abdel-Lateff, A., & Muller, D. (2006). Natural products from marine organisms and their associated microbes. *Chem Bio Chem*, 7, 229-238.

Krediet, C.J., Ritchie, K. B., Paul, V. J., & Teplitski, M. (2013). Coral-associated micro-organisms and their roles in promoting coral health and thwarting diseases. *Proceedings of the Royal Society B: Biological Sciences*, 280, 20122328.

Kumar, T. S., & Aparna, H. (2014). Anti-biofouling activity of Prodigiosin, a pigment extracted from *Serratia marcescens*. *Int. J. Curr. Microbiol. App. Sci*, 3(5), 712-725.

Kumar, V. S. (2012). *Population fluxes of commercially threatened sea cucumber species in Gulf of Mannar Biosphere Reserve for sustainable harvesting strategies to regularizing sea cucumber fishery*. A research report submitted to Man and

- Biosphere (MAB) Programme, UNESCO & ZSI, MoEnF, Government of India, New Delhi.
- Lampe, K. (2013). Holothurian density, distribution and diversity comparing sites with different degrees of exploitation in the shallow lagoons of Mauritius. *SPC Beche-de-mer Information Bulletin*, 33, 23-29.
- Lawrence, J. (1987). *A Functional Biology of Echinoderms*. Croom Helm, London & Sydney.
- Leal, M. C., Sheridan, C., Osinga, R., Dionísio, G., Rocha, R. J. M., Silva, B., Rosa, R. & Calado, R. (2014). Marine microorganism-invertebrate assemblages: Perspectives to solve the “supply problem” in the initial steps of drug discovery. *Marine drugs*, 12(7), 3929-3952.
- Leiva, S., Alvarado, P., Huang, Y., Wang, J., & Garrido, I. (2015). Diversity of pigmented Gram-positive bacteria associated with marine macroalgae from Antarctica. *FEMS microbiology letters*, 362(24).
- León-Palmero, E., Joglar, V., Álvarez, P. A., Martín-Platero, A., Llamas, I., & Reche, I. (2018). Diversity and antimicrobial potential in sea anemone and holothurian microbiomes. *PLoS ONE*, 13(5), e0196178.
- Lesson, R. P. (1830). *Centurie Zoologique On Choix D'animaux Rares, Nouveaux Ou Inparfaitement Connus*. Levrault, Paris, 244.
- Ley, R. E., Hamady, M., Lozupone, C., Turnbaugh, P. J., Ramey, R. R., Bircher, J. S., Schlegel, M. L., Tucker, T. A., Schrenzel, M. D., Knight, R., & Gordon, J. I. (2008). Evolution of mammals and their gut microbes. *Science (New York, N.Y.)*, 320(5883), 1647–1651.
- Long, R. A., & Azam, F. (2001). Antagonistic interactions among marine pelagic bacteria. *Applied and environmental microbiology*, 67(11), 4975–4983. <https://doi.org/10.1128/AEM.67.11.4975-4983.2001>
- Li, H., Qiao, G., Gu, J. Q., Zhou, W., Li, Q., Woo, S. H., Xu, D.H & Park, S. I. (2010). Phenotypic and genetic characterization of bacteria isolated from diseased cultured

- sea cucumber *Apostichopus japonicus* in northeastern China. *Diseases of aquatic organisms*, 91(3), 223-235.
- Li, J., Xu, Y., Jin, L., & Li, X. (2015). Effects of a probiotic mixture (*Bacillus subtilis* YB-1 and *Bacillus cereus* YB-2) on disease resistance and non-specific immunity of sea cucumber, *Apostichopus japonicus* (Selenka). *Aquaculture Research*, 46(12), 3008-3019.
- Li, J., Cao, J., Wang, X., Liu, N., Wang, W., & Luo, Y. (2017). *Acinetobacter pittii*, an emerging new multi-drug resistant fish pathogen isolated from diseased blunt snout bream (*Megalobrama amblycephala* Yih) in China. *Applied microbiology and biotechnology*, 101(16), 6459-6471.
- Li, J., & Engelberth, A. S. (2018). Quantification and purification of lutein and zeaxanthin recovered from distillers dried grains with solubles (DDGS). *Bioresources and Bioprocessing*, 5(1), 1-10.
- Li, Q., Qi, R. R., Wang, Y. N., Ye, S. G., Qiao, G., & Li, H. (2013). Comparison of cells free in coelomic and water-vascular system of sea cucumber, *Apostichopus japonicus*. *Fish & shellfish immunology*, 35(5), 1654-1657.
- Lima-Bittencourt, C. I., Cursino, L., Gonçalves-Dornelas, H., & Daniela, P. (2007). Multiple antimicrobial resistance in *Enterobacteriaceae* isolates from pristine freshwater. *Genetics and molecular research: GMR*, 6, 510-21.
- Lin, L. Y., Sulea, T., Szittner, R., Vassilyev, V., Purisima, E. O., & Meighan, E. A. (2001). Modeling of the bacterial luciferase-flavin mononucleotide complex combining flexible docking with structure-activity data. *Protein Science*, 10, 1563-1571.
- Liu, H., Tan, K. S., Zhang, X., Zhang, H., Cheng, D., Ting, Y., Li, S., Ma, H. & Zheng, H. (2020). Comparison of Gut Microbiota Between Golden and Brown Noble Scallop *Chlamys nobilis* and Its Association with Carotenoids. *Frontiers in microbiology*, 11, 36.
- Liu, Y., Lai, Q., Du, J., & Shao, Z. (2017). Genetic diversity and population structure of the *Bacillus cereus* group bacteria from diverse marine environments. *Scientific reports*, 7(1), 1-11.

- Long, R. A., & Azam, F. (2001). Antagonistic interactions among marine pelagic bacteria. *Applied and environmental microbiology*, 67(11), 4975-4983.
- Lu, Z., Yang, Q., Lin, Q., Wu, J., Du, H., & Zhou, C. (2021). Effects of *Bacillus* spp. on growth, gut microbiota compositions and immunity of sea cucumber (*Apostichopus japonicus*). *Aquaculture Research*.
- Lukman, A. L., Nordin, N. F. H., & Kamarudin, K. R. (2014). Microbial population in the coelomic fluid of *Stichopus chloronotus* and *Holothuria (Mertensiothuria) leucospilota* collected from Malaysian waters. *Sains Malaysiana*, 43(7), 1013-1021.
- MacTavish, T., Stenton-Dozey, J., Vopel, K., & Savage, C. (2012). Deposit-Feeding Sea Cucumbers Enhance Mineralization and Nutrient Cycling in Organically-Enriched Coastal Sediments. *PLoS ONE*, 7(11), e50031.
- Malik, K., Tokkas, J., & Goyal, S. (2012). Microbial Pigments: A review. *International Journal of Microbial Resource Technology*, 1(4), 361-365.
- Mani, P., Dineshkumar, G., Jayaseelan, T., Deepalakshmi, K., Kumar, C. G., & Balan, S. S. (2016). Antimicrobial activities of a promising glycolipid biosurfactant from a novel marine *Staphylococcus saprophyticus* SBPS 15. *3 Biotech*, 6(2), 1-9.
- Manon, M. V., Keerthana, G., & Preethi, K. (2015). Evaluation of antioxidant potential of bioactive colored metabolite isolated from *Exiguobacterium Profundum* BC2-11 and its bioactivities. *Int J Recent Sci Res*, 6(4), 3612-3617.
- Márta, D. (2012). *Molecular monitoring of meat spoiling Pseudomonas species analysis of Staphylococcal enterotoxin expression and formation* (Doctoral dissertation, Budapesti Corvinus Egyetem).
- Mashjoor, S., & Yousefzadi, M. (2019). Cytotoxic effects of three Persian Gulf species of Holothurians. *Iranian journal of veterinary research*, 20(1), 19.
- Mazlan, N., & Hashim, R. (2015). Spawning induction and larval rearing of the sea cucumber *Holothuria scabra* in Malaysia. *SPC Beche-de-mer Information Bulletin*, 35, 32-36.

Meddeb-Mouelhi, F., Moisan, J. K., Bergeron, J., Daoust, B., & Beaugard, M. (2016). Structural characterization of a novel antioxidant pigment produced by a photochromogenic *Microbacterium oxydans* strain. *Applied biochemistry and biotechnology*, 180(7), 1286-1300.

Mehmet, A., Huseyin, S., Bekir, T. Yilmaz, E., & Sevim, K. (2011). Proximate composition and fatty acid profile of three different fresh and dried commercial sea cucumber from Turkey. *International Journal of Food Science*, 46, 500-508.

Modesitt, D. E. (1970). Studies on predomination of selected bacteria of sewage origin. PhD thesis, Oklahoma State University.

Mohammadi, M., Burbank, L., & Roper, M. C. (2012). Biological role of pigment production for the bacterial phytopathogen *Pantoea stewartii* subsp. *stewartii*. *Applied and environmental microbiology*, 78(19), 6859-6865.

Morrow, K. M., Moss, A. G., Chadwick, N. E., & Liles, M. R. (2012). Bacterial associates of two Caribbean coral species reveal species-specific distribution and geographic variability. *Applied and environmental microbiology*, 78(18), 6438-6449.

Narsing Rao, M. P., Xiao, M., & Li, W. J. (2017). Fungal and bacterial pigments: secondary metabolites with wide applications. *Frontiers in microbiology*, 8, 1113.

Normand, P., Ponsonnet, C., Nesme, X., Neyra, M., & Simonet, P. (1996). ITS analysis of prokaryotes. *Molecular microbial ecology manual*, 1-12.

Nugraheni, S. A., Khoeri, M. M., Kusmita, L., Widyastuti, Y., & Radjasa, O. K. (2010). Characterization of carotenoid pigments from bacterial symbionts of seagrass *Thalassia hemprichii*. *J. Coast. Dev*, 14(1), 51-60.

Offret, C., Jégou, C., Mounier, J., Fleury, Y., & Le Chevalier, P. (2019). New insights into the haemo- and coelo-microbiota with antimicrobial activities from Echinodermata and Mollusca. *Journal of Applied Microbiology*, 126, 1023-1031.

O'Hara, A. M., & Shanahan, F. (2006). The gut flora as a forgotten organ. *EMBO Reports*, 7(7), 688-93.

- Ohfuji, K., Sato, N., Hamada-Sato, N., Kobayashi, T., Imada, C., Okuma, H., & Watanabe, E. (2004). Construction of a glucose sensor based on a screen-printed electrode and a novel mediator pyocyanin from *Pseudomonas aeruginosa*. *Biosensors and Bioelectronics*, 19(10), 1237-1244.
- Ojha, S., Kapoor, S., & Mishra, S. (2018). Carotenoid production by a novel isolate of *Microbacterium paraoxydans*. *Indian journal of microbiology*, 58(1), 118-122.
- Omran, N. E., & Allam, N. G. (2013). Screening of microbial contamination and antimicrobial activity of sea cucumber *Holothuria polii*. *Toxicology and Industrial Health*, 29(10), 944-54.
- Ong, S. H., Kukkillaya, V. U., Wilm, A., Lay, C., Ho, E. X. P., Low, L., Hibberd, M.L. & Nagarajan, N. (2013). Species identification and profiling of complex microbial communities using shotgun Illumina sequencing of 16S rRNA amplicon sequences. *PloS one*, 8(4), e60811.
- Osama, Y. A., Ridzwan, H., Muhammad, T., Jamaludin, M. D., Masa-Aki, I. & Zali, B. I. 2009. In vitro antioxidant and antiproliferative activities of three Malaysian sea cucumber species. *European Journal of Scientific Research*, 37(3): 376-387.
- Pagán-Jiménez, M., Ruiz-Calderón, J. F., Dominguez-Bello, M. G., & García-Arrarás, J. E. (2019). Characterization of the intestinal microbiota of the sea cucumber *Holothuria glaberrima*. *PloS one*, 14(1), e0208011.
- Pangestuti, R., & Arifin, Z. (2018). Medicinal and health benefit effects of functional sea cucumbers. *Journal of traditional and complementary medicine*, 8(3), 341-351.
- Pankor. (2004). Sea cucumber facts. Langkawi Magazine Nature Section. 16/08/2004. <http://www.best-of-langkawi.com/NATURE-gamut-problem.php>
- Petersen, L. E., Kellermann, M. Y., & Schupp, P. J. (2020). Secondary metabolites of marine microbes: From natural products chemistry to chemical ecology. *YOUMARES 9-The Oceans: Our Research, Our Future*, 159.

- Pelz, A., Wieland, K. P., Putzbach, K., Hentschel, P., Albert, K., & Götz, F. (2005). Structure and biosynthesis of staphyloxanthin from *Staphylococcus aureus*. *Journal of Biological Chemistry*, 280(37), 32493-32498.
- Piel, J. (2009). Metabolites from symbiotic bacteria. *Natural Product Reports*, 26, 338-362.
- Pierson, L. S., & Pierson, E. A. (2010). Metabolism and function of phenazines in bacteria: impacts on the behavior of bacteria in the environment and biotechnological processes. *Applied microbiology and biotechnology*, 86(6), 1659-1670.
- Pringgenies, D., Girsang, P. H., Yudiati, E., & Santosa, G. W. (2020). Exploration of sea cucumber intestinal symbiont microbe as probiotic microbe candidate in healthcare products. *JFMR (Journal of Fisheries and Marine Research)*, 4(1), 27-34.
- Priyaja, P., Rosamma, P., & Bright Singh, I. S. (2012). *Pyocyanin (5-methyl-1-hydroxyphenazine) produced by Pseudomonas aeruginosa as antagonist to vibrios in aquaculture: Overexpression, downstream process and toxicity* (Doctoral dissertation, Cochin University of Science and Technology).
- Proksch, P., Edrada, R., & Ebel, R. (2002). Drugs from the seas—current status and microbiological implications. *Applied microbiology and biotechnology*, 59(2), 125-134.
- Purcell, S., Samyn, Y., & Conand, C. (2012). Commercially important sea cucumbers of the world. FAO Species Catalogue for Fishery Purposes. No 6. Rome, FAO. 150pp.
- Radjasa, O. K., & Sabdono, A. (2009). Bacterial symbionts of reef's invertebrates as a sustainable source of marine natural products. *Current Research in Bacteriology*, 2(1), 7-13.
- Raguénes, G., Moppert, X., Richert, L., Ratiskol, J., Payri, C., Costa, B., & Guezennec, J. (2004). A novel exopolymer-producing bacterium, *Paracoccus zeaxanthinifaciens* subsp. *payriae*, isolated from a “kopara” mat located in Rangiroa, an atoll of French Polynesia. *Current microbiology*, 49(3), 145-151.

Rahman, M. A., Yusoff, F. M., & Arshad, A. (2015). Sea cucumber fisheries: Global status, culture, management and extinction risks. *International Journal of Chemical, Environmental and Biological Sciences*, 3(4), 344-348.

Rahman, M. A. (2014). Culture potentials of sea cucumbers (Echinodermata: Holothuroidea) and their biomedical applications. *International Conference on Chemical, biological and Environmental Sciences*. Kuala Lumpur, Malaysia. May 12-13, 2014.

Ramesh, C., Vinithkumar, N. V., & Kirubakaran, R. (2019). Marine pigmented bacteria: A prospective source of antibacterial compounds. *Journal of Natural Science, Biology and Medicine*, 10(2), 104.

Ramírez-González, J., Moity, N., Andrade-Vera, S., & Reyes, H. (2020). Overexploitation and more than a decade of failed management leads to no recovery of the Galapagos sea cucumber fishery. *Frontiers in Marine Science*, 7, 920.

Rasko, D. A., Altherr, M. R., Han, C. S., & Ravel, J. (2005). Genomics of the *Bacillus cereus* group of organisms. *FEMS microbiology reviews*, 29(2), 303-329.

Reef Check Malaysia. (2018). Status of coral reefs in Malaysia, 2015. *Reef check malaysia survey report* Department of Marine Park, Malaysia.

Rehan, M. B. M. (2012). *Multiple mechanisms mediating the starvation induced activation of recombination at HIS4 in Saccharomyces cerevisiae* (Doctoral dissertation, University of Leicester).

Ridzwan, B. H., Kaswandi, M. A., Azman, Y., & Fuad, M. (1995). Screening for antibacterial agents in three species of sea cucumbers from coastal areas of Sabah. *General Pharmacology*, 26(7), 1539-1543.

Ridzwan, B. H. (2007). Sea cucumbers: a Malaysian heritage. *Research Centre of International Islamic University Malaysia (IIUM), Kuala Lumpur Wilayah Persekutuan: Kuala Lumpur, Malaysia*.

Rinehart, K. (2000). Antitumor Compounds from Tunicates. *Medicinal research reviews*, 20, 1-27.

Roberts, D., Moore, H. M., Berges, J., Patching, J. W., & Carton, M. W. (2001). Sediment distribution, hydrolytic enzyme profiles and bacterial activities in the guts of *Oneirophanta mutabilis*, *Psychropotes longicauda* and *Pseudostichopus villosus*: what do they tell us about digestive strategies of abyssal holothurians? *Progress in Oceanography*, 50, 443–458.

Rodriguez-Amaya, D. B. (1999). A Guide to Carotenoid Analysis in Foods. ILSI Press, Washington, DC., pp: 14-15.

Rohwer, F., Seguritan, V., Azam, F., & Knowlton, N. (2002). Diversity and distribution of coral-associated bacteria. *Marine Ecology Progress Series*, 243, 1-10.

Romalde, J. L., Diéguez, A. L., Lasa, A., & Balboa, S. (2014). New *Vibrio* species associated to molluscan microbiota: a review. *Frontiers in Microbiology*, 4, 413.

Saitou, N., & Nei, M. (1987). The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, 4, 406–425.

Sajilata, M. G., Singhal, R. S., & Kamat, M. Y. (2008). The carotenoid pigment zeaxanthin—a review. *Comprehensive reviews in food science and food safety*, 7(1), 29-49.

Sasidharan, P., Raja, R., Karthik, C., Ranandkumar, S., & Indra Arulselvi, P. (2013). Isolation and characterization of yellow pigment producing *Exiguobacterium* sps.

Savini, V., Catavittello, C., Carlino, D., Bianco, A., Pompilio, A., Balbinot, A., Piccolomini, R., Di Bonaventura, G. & D'Antonio, D. (2009). *Staphylococcus pasteurii* bacteraemia in a patient with leukaemia. *Journal of clinical pathology*, 62(10), 957-958.

Schneider, K. L., Marrero, G., Alvarez, A. M., & Presting, G. G. (2011). Classification of Plant Associated Bacteria Using RIF, a Computationally Derived DNA Marker. *PLoS ONE*, 6(4).

- Schneider, K., Silverman, J., Woolsey, E., Eriksson, H., Byrne, M., & Caldeira, K. (2011). Potential influence of sea cucumbers on coral reef CaCO₃ budget: A case study at One Tree Reef. *Journal of Geophysical Research: Biogeosciences*, 116(G4).
- Schwartzmann, G., da Rocha, A.B., Berlinck, J.G.S., & Jimeno, J. (2001). Marine organisms as a source of new anticancer agents. *The Lancet Oncology*, 2, 221– 225.
- Semper, C. (1868). Reisen im Archipel der Philippinen. 2. Wissenschaftliche Resultate. 1. Holothurien. Wilhelm Engelmann, Leipzig.
- Sha, Y., Liu, M., Wang, B., Jiang, K., Sun, G., & Wang, L. (2016). Gut bacterial diversity of farmed sea cucumbers *Apostichopus japonicus* with different growth rates. *Microbiology*, 85(1), 109-115.
- Shakouri, A., Nabavi, M. B., Kochanian, P., Savari, A., Safahieh, A., & Aminrad, T. (2009). New observation of three species of sea cucumbers from Chabahar Bay (Southeast coasts of Iran). *Journal of Biological Sciences*, 9(2), 184-187.
- Sharif, A., & Osman, H. (2016, February 16). Prospek cerah pulihara gamat. *Harian Metro*. Ruang Setempat (p. 34). Langkawi, Malaysia.
- Shawkey, M. D., & D'Alba, L. (2017). Interactions between colour-producing mechanisms and their effects on the integumentary colour palette. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 372(1724), 20160536.
- Shi, W., Zhang, J., Wang, Y., Ji, J., Guo, L., Ren, Y., Qiao, G., Wang, Q. & Li, Q. (2020). Transcriptome analysis of sea cucumber (*Apostichopus japonicus*) polian vesicles in response to evisceration. *Fish & shellfish immunology*, 97, 108-113.
- Siahaan, E. A., Pangestuti, R., Munandar, H., & Kim, S-K. (2017). Cosmeceuticals Properties of Sea Cucumbers: Prospects and Trends. *Cosmetics*, 4(3), 26.
- Silhavy, T., Kahne, D., & Walker, S. (2010). The Bacterial Cell Envelope. *Cold Spring Harbor perspectives in biology*, 2(5).
- Soliev, A. B., Hosokawa, K., & Enomoto, K. (2011). Bioactive pigments from marine bacteria: applications and physiological roles. *Evidence-based complementary and alternative medicine: eCAM*, 2011.

Staley, J. T., Castenholz, R. W., Colwell, R. R., Holt, J. G., Kane, M. D., Pace, N. R., Salyers, A. A., & Tiedje, J. M. (1997). The Microbial World: Foundation of the Biosphere. The American Academy of Microbiology. The American Academy of Microbiology, Washington, DC. Online, <http://www.asmtusa.org/acasrc/aca1/html>.

Sulardiono, B., Widyorini, N., Suprpto, D., Ayuningrum, D., & Rahman, A. (2020). Evaluation of antibacterial activity and molecular characterization of bacteria from *Holothuria atra* intestine collected from anthropogenic and non-anthropogenic region in Karimunjawa, Indonesia. *Biodiversitas Journal of Biological Diversity*, 21(7).

Sung, C., Kim, B. G., Kim, S., Joo, H. S., & Kim, P. I. (2010). Probiotic potential of *Staphylococcus hominis* MBBL 2–9 as anti-*Staphylococcus aureus* agent isolated from the vaginal microbiota of a healthy woman. *Journal of applied microbiology*, 108(3), 908-916.

Starr, M. P. (1958). The blue pigment of *Corynebacterium insidiosum*. *Archives of Microbiology*, 30: 325-334.

Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., & Kumar, S. (2011). MEGA5: Molecular Evolutionary Genetics Analysis Using Maximum Likelihood, Evolutionary Distance, And Maximum Parsimony Methods. *Molecular biology and evolution*, 28, 2731-9.

Tangestani, M., & Kunzmann, A. (2019). Isolation and characterization of bacteria from the lesion of juvenile sea cucumber *Holothuria scabra* (Jaeger, 1938) with symptom of skin ulceration disease. *Iranian Journal of Fisheries Sciences*, 18(4), 915-923.

Taylor, M. W., Schupp, P. J., De Nys, R., Kjelleberg, S., & Steinberg, P. D. (2005). Biogeography of bacteria associated with the marine sponge *Cymbastela concentrica*. *Environmental microbiology*, 7(3), 419-433.

Tehranifard, A., & Rahimibashar, M. R. (2012). Description of a sea cucumber species *Holothuria atra* Jaeger, 1833 from Kish Island Iran (Echinodermata: Holothuroidea). *Journal of Basic and Applied Scientific Research*, 2, 12660-12664.

Theis, K.R., Dheilly, N. M., Klassen, J. L., Brucker, R. M., Baines, J. F., Bosch, T. C., Cryan, J. F., Gilbert, S. F., Goodnight, C. J., Lloyd, E. A., Sapp, J., Vandenkoornhuys, P., Zilber-Rosenberg, I., Rosenberg, E., & Bordenstein, S. R. (2016). Getting the Hologenome Concept Right: an Eco-Evolutionary Framework for Hosts and Their Microbiomes. *mSystems*, 1(2):e00028-16.

Thompson, J. D., Higgins, D. G., & Gibson, T. J. (1994). *Computer Application in the Biosciences*, 10, 19–29.

Tuli, H. S., Chaudhary, P., Beniwal, V., & Sharma, A. K. (2015). Microbial pigments as natural color sources: current trends and future perspectives. *Journal of food science and technology*, 52(8), 4669–4678.

Usman, H. M., Abdulkadir, N., Gani, M., & Maiturare, H. (2017). Bacterial pigments and its significance. *MOJ Bioequivalence and Bioavailability*, 4, 285–288.

Uticke, S., Purcell, S. W., & Blockmans, B. (2005). Natural hybridization does not dissolve species boundaries in commercially important sea cucumbers. *Biological Journal of the Linnean Society*, 85, 261-270.

VandenSpiegel, D., & Jangoux, M. (1987). Cuvierian tubules of the holothuroid *Holothuria forskali* (Echinodermata): a morphofunctional study. *Marine Biology*, 96(2), 263-275.

Velho-Pereira, S., & Furtado, I. (2012). Antibacterial activity of halophilic bacterial bionts from marine invertebrates of Mandapam-India. *Indian Journal of Pharmaceutical Sciences*, 74(4), 331.

Velmurugan, P., Venil, C. K., Veera Ravi, A., & Dufossé, L. (2020). Marine bacteria are the cell factory to produce bioactive pigments: A prospective pigment source in the ocean. *Frontiers in Sustainable Food Systems*, 4, 234.

Venil, C. K., Zakaria, Z. A. & Ahmad, W. A. (2013). Bacterial pigments and their applications. *Process Biochemistry*, 48, 1065-1079.

- Walford J. (ed.) (1980). "Historical development of food colouration," in *Developments in Food Colours*, (London: Applied Science publishers;)
- Wang Y, Sheng H-F, He Y, Wu J-Y, Jiang Y-X, Tam NF-Y, Zhou H-W. (2012) Comparison of the levels of bacterial diversity in freshwater, intertidal wetland, and marine sediments by using millions of illumina tags. *Appl Environ Microbiol.*, 78,8264-8271.
- Ward-Rainey, N., Rainey, F. A., & Stackebrandt, E. (1996). A study of the bacterial flora associated with *Holothuria atra*. *Journal of Experimental Marine Biology and Ecology*, 203, 11-26.
- Weisburg, W. G., Barns, S. M., Pelletier, D. A., & Lane, D. J. (1991). 16S ribosomal DNA amplification for phylogenetic study. *Journal of Bacteriology*, 173, 697-703.
- Wibowo, J. T., Kellermann, M. Y., Versluis, D., Putra, M. Y., Murniasih, T., Mohr, K. I., Wink, J., Engelmann, M., Praditya, D.F., Steinmann, E. & Schupp, P. J. (2019). Biotechnological potential of bacteria isolated from the sea cucumber *Holothuria leucospilota* and *Stichopus vastus* from Lampung, Indonesia. *Marine drugs*, 17(11), 635.
- Wiese, J., & Imhoff, J. F. (2019). Marine bacteria and fungi as promising source for new antibiotics. *Drug development research*, 80(1), 24-27.
- Wijaya, A. P., Bondar, K. G., Frederick, E. H., Igarashi, Y., & Sibero, M. T. (2020, October). Identification of marine bacteria HPP. 4A and HPP. T13 and its anticancer activity against P388 murine leukaemia cell. In *IOP Conference Series: Earth and Environmental Science* (Vol. 584, No. 1, p. 012005). IOP Publishing.
- Wilson, G. S., Raftos, D. A., Corrigan, S. L., & Nair, S. V. (2010). Diversity and antimicrobial activities of surface-attached marine bacteria from Sydney Harbour, Australia. *Microbiological research*, 165(4), 300-311.
- Woese, C. R. (1987). Bacterial evolution. *Microbiological reviews*, 51(2), 221-271.

- Wong, D., Nielsen, T. B., Bonomo, R. A., Pantapalangkoor, P., Luna, B., & Spellberg, B. (2017). Clinical and pathophysiological overview of Acinetobacter infections: a century of challenges. *Clinical microbiology reviews*, 30(1), 409-447.
- Xu, L. Q., Zeng, J. W., Jiang, C. H., Wang, H., Li, Y. Z., Wen, W. H., Li, J.H., Wang, F., Ting, W.J., Sun, Z.Y. & Huang, C. Y. (2017). Isolation and determination of four potential antimicrobial components from *Pseudomonas aeruginosa* extracts. *International journal of medical sciences*, 14(13), 1368.
- Xu, H., Wang, L., Bao, X., Jiang, N., Yang, X., Hao, Z., Chang, Y. & Ding, J. (2019). Microbial communities in sea cucumber (*Apostichopus japonicus*) culture pond and the effects of environmental factors. *Aquaculture Research*, 50(4), 1257-1268.
- Yan, F. J., Tian, X. L., Dong, S. L., Fang, Z. H., & Yang, G. (2014). Growth performance, immune response, and disease resistance against *Vibrio splendidus* infection in juvenile sea cucumber *Apostichopus japonicus* fed a supplementary diet of the potential probiotic *Paracoccus marcusii* DB11. *Aquaculture*, 420-421, 105-111.
- Yang, H. (Ed.). (2015). *The sea cucumber Apostichopus japonicus: history, biology and aquaculture*. Academic Press.
- Yasoda, H. N., Chi, Z., & Ling, Z. K. (2006). Probiotics and sea cucumber farming. *SPC Beche-de-mer Information Bulletin*, 24, 45-48.
- Yolmeh, M. (2016). Antimicrobial activity of pigments extracted from *Rhodotorula glutinis* against some bacteria and fungi.
- Zaghdoudi, K., Ngomo, O., Vanderesse, R., Arnoux, P., Myrzakhmetov, B., Frochot, C., & Guiavarc'h, Y. (2017). Extraction, identification and photo-physical characterization of persimmon (*Diospyros kaki* L.) carotenoids. *Foods*, 6(1), 4.
- Zhang, X., Nakahara, T., Miyazaki, M., Nogi, Y., Taniyama, S., Arakawa, O., Inoue, T. & Kudo, T. (2012). Diversity and function of aerobic culturable bacteria in the intestine of the sea cucumber *Holothuria leucospilota*. *The Journal of General and Applied Microbiology*, 58(6), 447-456.

Zhang G., Zhang Y., Yin X., & Wang S. (2015). *Nesterenkonia alkaliphila* sp. nov., an alkaliphilic, halotolerant actinobacteria isolated from the western Pacific Ocean. *International Journal of Systematic and Evolutionary Microbiology*, 65, 516–521.

Zhang, Y., Liu, Z., Sun, J., Xue, C., & Mao, X. (2018). Biotechnological production of zeaxanthin by microorganisms. *Trends in Food Science & Technology*, 71, 225-234.

Zhang, H., Wang, Q., Liu, S., Huo, D., Zhao, J., Zhang, L., Zhao, Y., Sun, L. & Yang, H. (2019). Genomic and metagenomic insights into the microbial community in the regenerating intestine of the sea cucumber *Apostichopus japonicus*. *Frontiers in microbiology*, 10, 1165.

Zinger, L., Amaral-Zettler, L. A., Fuhrman, J. A., Horner-Devine, M. C., Huse, S. M., Welch, D. B. M., Martiny, J.B., Sogin, M., Boetius, A. & Ramette, A. (2011). Global patterns of bacterial beta-diversity in seafloor and seawater ecosystems. *PloS one*, 6(9), e24570.

Zulfaqar, S., Rahman, M. A., & Yusoff, F. M. (2016). Trends, prospects and utilizations of sea cucumber fisheries in Malaysia. *Int J Adv Agric Environ Engg*, 3(1), 114-116.

Zulkifli, Y., Alitheen, N. B., Son, R., Yeap, S. K., Lesley, M. B., & Raha, A. R. (2009). Identification of *Vibrio parahaemolyticus* isolates by PCR targeted to the *toxR* gene and detection of virulence genes. *International Food Research Journal*, 16, 289-296.